


AGNAR[®]

(Calcium + Vitamin D₃ + Vitamin K₂)
Film-coated Tablets
500mg + 400IU + 90mcg



DESCRIPTION

Agnar Tablet is a nutritional supplement for bone care and maintenance of good health having a combination of Calcium, Vitamin D₃ and Vitamin K₂. Agnar is used to prevent or treat low blood calcium levels in people who do not get enough calcium from their diet. Vitamin D helps your body absorb calcium and phosphorus and is vital to bone formation, blood pressure regulation, hormone production, immune and nervous system function. Vitamin K is very effective in bone and heart health.

COMPOSITION

Agnar film-coated tablet contains:
Vitamin D₃ (Cholecalciferol) Food Grade... 400IU
Vitamin K₂ (MK-7) Food Grade...90mcg
Elemental Calcium... 500mg

INDICATIONS

Agnar Tablets are used to prevent and/or treat conditions related to Calcium, Vitamin D and Vitamin K deficiency such as:

- Bone loss (Osteoporosis)
- Weak bone (Osteomalacia)
- Rickets in children
- Decreased activity of parathyroid gland (Hypoparathyroidism)

Agnar Tablets maintain and support:

- Optimal bone health and heart health
- Recommended dietary intake of calcium & vitamin D₃ for pregnant, nursing & postmenopausal women
- Healthy active lifestyle

DOSAGE AND ADMINISTRATION

Take Agnar 1-2 tablets daily with food or as directed by the healthcare practitioner.

ADVERSE EFFECTS

Agnar Tablets may cause constipation, increased calcium levels in blood or urine, abdominal pain, stomach upset, nausea, vomiting, loss of appetite, unusual weight loss, mental/mood changes, bone/muscle pain, headache, increased thirst and urination, weakness, tiredness, fast/pounding heartbeat and serious allergic reactions including rash, itching/swelling, severe dizziness, trouble breathing.

CONTRAINDICATIONS

Agnar Tablets must not be used in patients:

- With hypersensitivity to any of the ingredient of the product.
- Who are on warfarin therapy.
- With increased levels of calcium in blood or urine.
- With kidney stones.
- With severe renal impairment and renal failure.
- With increased levels of Vitamin D.

PRECAUTIONS

- Caution is advised in patients who have serious medical condition and who are taking other medications.
- During long-term treatment, serum calcium levels should be followed and renal function should be monitored through measurements of serum creatinine.

PREGNANCY & NURSING WOMEN

Agnar Tablets may be given during pregnancy and nursing. However, pregnant and nursing women should consult a healthcare practitioner prior taking Agnar Tablets.

OVERDOSAGE

Overdose can lead to hypervitaminosis and hypercalcaemia. Persistently high calcium levels may lead to irreversible renal damage and soft tissue calcification.

Treatment: Use of Agnar Tablets must be discontinued. Serum electrolytes, renal function and diuresis must be monitored.

STORAGE:

Store below 30°C.
Protect from sunlight and moisture.
Keep out of reach of children.

PRESENTATION

Agnar film-coated tablets are available in blister pack of 30's.

This is not an Allopathic product.

Manufactured for:



29-30/27, K.I.A., Karachi, Pakistan

By: Novamed Healthcare (Pvt) Ltd.,
28-km Ferozepur Road, Lahore, Pakistan